

Level-Rank Duality in Kazama-Suzuki Models

Tibra Ali†

D.A.M.T.P., C.M.S., Wilberforce Road, Cambridge University, Cambridge CB3 0WA, England
(preprint DAMTP-2002-11; hep-th/0201214)

We give a path-integral proof of level-rank duality in Kazama-Suzuki models for world-sheets of spherical topology.

Since the discovery of nonabelian bosonization in two dimensions [1,2] we have learned quite a bit about the conformal structure of string theory and two dimensional statistical field theory. An important class of supersymmetric conformal field theories was discovered by Kazama and Suzuki [KS] [3]. These models are believed to be that subset of the $N = 1$ supersymmetric Goddard-Kent-Olive [GKO] [4] coset models which admit $N = 2$ superconformal symmetry. Therefore, they give explicit integrable conformal field theory realization of target spaces with space-time supersymmetry. Since in a lagrangian formulation coset models correspond to the gauging a WZW model by $g(z, \bar{z}) \rightarrow h(z, \bar{z})g(z, \bar{z})h^{-1}(z, \bar{z})$ they describe, in string theory, space(times) with black holes [5–7]. Some of the KS models have been conjectured to enjoy the following duality property [3]:

$$\begin{aligned} & [SU(M+N)/(SU(M) \times SU(N) \times U(1))]_k \\ & \approx [SU(k+M)/(SU(M) \times SU(k) \times U(1))]_N \end{aligned} \quad (1)$$

and similarly

$$\begin{aligned} & [SO(N+2)/(SO(N) \times SO(2))]_k \\ & \approx [SO(k+2)/(SO(k) \times SO(2))]_N. \end{aligned} \quad (2)$$

There is now a considerable body of work on this duality, especially in the operator approach [8,9] but a path-integral formulation of the duality has been lacking. It is desirable for aesthetic as well as string theoretic reasons to have a path-integral analysis of this intriguing duality. The purpose of this letter is to supply such a proof for world-sheet of spherical topology. We shall give our results for the case $[SU(N+1)/(SU(N) \times U(1))]_k \approx [SU(k+1)/(SU(k) \times U(1))]_N$ in detail but the method can be generalized for all the models involved.

It has been known for some time that an $SU(N)$ WZW model at level k describes N Dirac *parafermions* of order k [10,11]:

$$\mathcal{Z}_1 = \int [dg] e^{-kI_{SU(N)}(g)} = \int [d\psi][d\bar{\psi}][dA] e^{-I_F(\psi, A)} \quad (3)$$

where $I_{SU(N)}$ is the level-1 WZW action for the group $SU(N)$ and the fermionic action is given by $I_F(\psi, A) = \frac{1}{2\pi} \int d^2z i\bar{\psi}_a^i \not{\partial} \psi_a^i + \bar{\psi}_a^i A^I \lambda_{ij}^I \psi_a^j$ with $i, j = 1, \dots, k$, $I = 1, \dots, k^2$ and $a = 1, \dots, N$. Because of its equivalence to the constrained fermionic system the $SU(N)$ WZW model at level k can also be expressed in terms of a negative level WZW model, free fermions, a free boson and ghosts [12]:

$$\begin{aligned} \mathcal{Z}_1 &= \int [d\bar{\psi}][d\psi][d\hat{g}][d\Phi] \det \partial_+ \det \partial_- e^{-I_B(\Phi)} e^{-I_F(\psi, 0)} \\ &\times e^{-[-N-2k]I_{SU(k)}(\hat{g})}. \end{aligned} \quad (4)$$

Where $I_B(\Phi)$ is the action of a free boson. Negative level WZW models are non-unitary but since they arise here with ghosts, it is believed that a BRST cohomology structure ensures their unitarity. This point has been explored in [13].

There are various approaches to obtaining a supersymmetric extension of a gauged-WZW model on G/H [14–16]. The one that we follow here is the one by Witten [14]. In this approach the supersymmetric gauged-WZW action is given by

$$\begin{aligned} I(g, A, \psi) &= I(g, A) \\ &+ \frac{i}{4\pi} \int d^2z \text{tr}(\psi_+ D_- \psi_+ + \psi_- D_+ \psi_-). \end{aligned} \quad (5)$$

Here $I(g, A)$ is the bosonic action for a gauged-WZW model and the $\psi_{\pm}$ are Weyl fermions which belong to the complexified orthogonal complement of the Lie algebra $\mathcal{H}$ and they are minimally coupled to the gauge fields $A_{\pm}$. The condition that the $N = 1$ superconformal algebra can be extended to $N = 2$ superconformal algebra can be translated to the condition that the group coset G/H is Kähler. Eq. (5) can then be written in a suitable form to reflect that structure [14].

We are interested in the partition function for the KS models as gauged-WZW models:

$$\begin{aligned} \mathcal{Z}_2 &= \int [dg][dA_+][dA_-][d\psi_+][d\psi_-] e^{-kI(g, A, \psi)} \\ &= \int [dg][dh][d\tilde{h}][d\psi_+][d\psi_-] \det D_+ \det D_- \\ &\times e^{-kI_G(g)} e^{kI_H(h^{-1}\tilde{h})} e^{\frac{1}{2}(c_G - c_H)I_H(h^{-1}\tilde{h})} e^{I_F(\psi, 0)}. \end{aligned} \quad (6)$$

Where we have made the change of variables $A_- = \partial_- \tilde{h} h^{-1}$, $A_+ = \partial_+ h h^{-1}$ [13]. c_G (c_H) is the quadratic Casimir operator of the group G (H) in the adjoint representation. In the above expression the level $-\frac{1}{2}(c_G - c_H)$ WZW model arises from “rotating away” the gauge coupling to the fermions [16]. We can now use the following identity [2]

$$\det D_+ \det D_- = e^{c_H I_H(h^{-1}\tilde{h})} \det \partial_+ \det \partial_- \quad (7)$$

and rewrite eq. (6) in a gauge fixed form (which does not introduce any new ghosts):

$$\mathcal{Z}_2 = \int [dg][dh][d\psi_+][d\psi_-] \det \partial_+ \det \partial_- \times e^{-kI_G(g)} e^{(k+\frac{1}{2})(c_G-c_H)+c_H} I_H(h) e^{-I_F(\psi,0)}. \quad (8)$$

We now specialize to the case where $G = SU(N+1)$ and $H = SU(N) \times U(1)$. Using eqs. (3) & (4) this can be written as

$$\begin{aligned} \mathcal{Z}_2 &= \int [d\hat{g}][d\psi'_-][d\psi'_+][d\Phi'] \det \partial'_+ \det \partial'_- \\ &\times e^{-I_B(\Phi')} e^{-[-N-1-2k]I_{SU(k)}(\hat{g})} e^{-I_F(\psi',0)} \\ &\times [dh][d\Phi][d\psi_+][d\psi_-] \det \partial_+ \det \partial_- \\ &\times e^{-I_B(\Phi)} e^{(k+1+2N)I_{SU(N)}(h)} e^{-I_F(\psi,0)}. \end{aligned} \quad (9)$$

In the above expression primed spinors correspond to free Dirac spinors with global $SU(N+1) \times SU(k) \times U(1)$ symmetry. The $Nk+k$ free Dirac spinors ψ' can be combined with the original $2N$ Weyl spinors ψ to give $Nk+N$ free Dirac spinors χ' and $2k$ free Weyl spinors χ . Having rearranged the spinors in this way eqs. (3) & (4) can be applied to the level $-(k+1+2N)$ WZW model on $SU(N) \times U(1)$ to yield

$$\begin{aligned} \mathcal{Z}_2 &= \int [d\hat{h}][d\hat{g}][d\chi_+][d\chi_-][d\Phi] \det \partial'_+ \det \partial'_- \\ &\times e^{-I_B(\Phi)} e^{-NI_{SU(k+1)}(\hat{h})} \\ &\times e^{-[-N-1-2k]I_{SU(k)}(\hat{g})} e^{-I_F(\chi,0)}. \end{aligned} \quad (10)$$

This is the gauge-fixed partition function for a $[SU(k+1)/(SU(k) \times U(1))]_N$ gauged-WZW model with $\hat{h} \in SU(k+1)$ and $\hat{g} \in SU(k)$. The central charge can be easily obtained from eq.(9) which is symmetric in k & N . In an obvious notation the central charge is given by

$$\begin{aligned} C_{\text{TOT}} &= C(SU(k), -N-1-2k) \\ &+ C(SU(N), -k-1-2N) \\ &+ C'_{ghost} + C_{ghost} + 2C_\Phi + C_{\psi'} + C_\psi \end{aligned} \quad (11)$$

where $C(G, k) = \frac{2k \dim(G)}{2k+c_G}$ is the standard central charge for a level- k WZW model on G , $C'_{ghost} = -2k^2$, $C_{ghost} = -2N^2$, $C_\Phi = 1$, $C_{\psi'} = Nk+k$ and $C_\psi = N$, which give

$$C_{\text{TOT}} = \frac{3kN}{k+N+1} \quad (12)$$

which is the anomaly found by KS [3].

The above proof can be extended easily to all the models involved in (1). Following the same set of steps as before we arrive at an expression for the partition function

$$\begin{aligned} \mathcal{Z}_3 &= \int [d\hat{g}][dh_1][dh_2][d\psi'_+][d\psi'_-][d\psi'_+][d\psi'_-][d\Phi][d\Phi'] \\ &\times e^{(M+N+2k)I_{SU(k)}(\hat{g})} e^{-I_B(\Phi')} \det \partial'_+ \det \partial'_- \\ &\times e^{(k+N+2M)I_{SU(M)}(h_1)} \det \partial_{1+} \det \partial_{1-} \\ &\times e^{(k+M+2N)I_{SU(N)}(h_2)} e^{-I_B(\Phi)} \det \partial_{2+} \det \partial_{2-} \\ &\times e^{-I_F(\psi',0)} e^{-I_F(\psi,0)} \end{aligned} \quad (13)$$

where we have written the ghosts on the same line as the corresponding WZW model. This expression is completely symmetric in k , M and N and it can be easily verified that conformal anomaly agrees with [3]. By using analogous identities [17] for orthogonal groups this analysis may be extended to (2).

Eqs. (9) & (13) lay bare the structure of the level-rank duality in a concise form. It can be used to study the precise map between the correlation functions of the dual theories, modular invariance and coupling to gravity. Also interesting is the fact that this duality connects a string theory vacuum in the semiclassical limit (for k , N or $M \rightarrow \infty$) to a string theory without a geometric interpretation (and with infinite naive “dimensions”). These issues are under investigation [18].

ACKNOWLEDGEMENTS

It is a pleasure to thank Malcolm Perry & Adam Ritz for helpful discussions. I am also grateful to Jürgen Fuchs & Christoph Schweigert for pointing out additional references and insightful comments for further investigations.

[†]E-Mail: T.Ali@damtp.cam.ac.uk

-
- [1] E. Witten, Commun. Math. Phys. **92** (1984) 455.
 - [2] A. M. Polyakov and P. B. Wiegmann, Phys. Lett. B **131** (1983) 121. Phys. Lett. B **141** (1984) 223.
 - [3] Y. Kazama and H. Suzuki, Nucl. Phys. B **321** (1989) 232.
 - [4] P. Goddard, A. Kent and D. I. Olive, Phys. Lett. B **152** (1985) 88. Commun. Math. Phys. **103** (1986) 105.
 - [5] E. Witten, Phys. Rev. D **44** (1991) 314.
 - [6] A. R. Lugo, Phys. Rev. D **52** (1995) 2266
 - [7] K. Bardacki, M. J. Crescimanno and E. Rabinovici, Nucl. Phys. B **344** (1990) 344.
 - [8] S. G. Naculich and H. J. Schnitzer, Nucl. Phys. B **505** (1997) 727
 - [9] J. Fuchs and C. Schweigert, Annals Phys. **234** (1994) 102
 - [10] I. Antoniadis and C. Bachas, Nucl. Phys. B **278** (1986) 343.
 - [11] A. N. Redlich and H. J. Schnitzer, Phys. Lett. B **167** (1986) 315 [Erratum-ibid. B **193** (1986) 536].
 - [12] S. G. Naculich and H. J. Schnitzer, Nucl. Phys. B **347** (1990) 687.
 - [13] D. Karabali and H. J. Schnitzer, Nucl. Phys. B **329** (1990) 649.
 - [14] E. Witten, Nucl. Phys. B **371** (1992) 191.
 - [15] H. J. Schnitzer, Nucl. Phys. B **324** (1989) 412.
 - [16] A. A. Tseytlin, Nucl. Phys. B **411** (1994) 509
 - [17] A. N. Redlich and H. J. Schnitzer, Phys. Lett. B **193** (1987) 471.
 - [18] T. Ali *in preparation*.